

# Nitromusk and Polycyclic Musk Compounds as Long-term Inhibitors of Cellular Xenobiotic Defense Systems Mediated by Multi-Drug Transporters

Till Luckenbach and David Epel

doi:10.1289/ehp.7301 (available at <http://dx.doi.org/>)

Online 30 September 2004



**Nitromusk and Polycyclic Musk Compounds as Long-term  
Inhibitors of Cellular Xenobiotic Defense Systems Mediated by  
Multi-Drug Transporters**

Till Luckenbach and David Epel

Hopkins Marine Station of Stanford University, 120 Oceanview Blvd

Pacific Grove, CA 93950, USA

Corresponding author/express mail address

David Epel

Hopkins Marine Station of Stanford University

120 Oceanview Blvd

Pacific Grove, CA 93950

USA

T: 831-655 6226

F: 831-375 0793

E: [depel@stanford.edu](mailto:depel@stanford.edu)

## **Running title**

Synthetic musk compounds as MXR inhibitors

## **Keywords**

nitromusks, polycyclic musks, fragrances, chemosensitizers, multixenobiotic resistance, MXR, multidrug resistance, MDR, *Mytilus californianus*

## **Acknowledgements**

This publication was supported in part by the German Academic Exchange Service (DAAD), in part by the National Sea Grant College Program of the U.S. Department of Commerce's National Oceanic and Atmospheric Administration under Grant # R/CZ-182 through the California Sea Grant College Program; and in part by the California State Resources Agency. The views expressed herein do not necessarily reflect the views of any of those organizations. The authors would like to thank Shobha Sadasivaiah for technical assistance, Jim Watanabe for help with the statistical analyses, Ilaria Corsi, Amro Hamdoun, Tvrtko Smital and Cathy Thaler for discussions and comments on the manuscript. The authors declare they have no conflict of interest.

## **Abbreviations**

FSW: filtered seawater,  $K_{ow}$ : octanol-water coefficient, MXR: Multixenobiotic resistance, PPCPs: pharmaceutical and personal care products, MK: musk ketone, MX: musk xylene, for abbreviations of other compounds see Table 1

## **Section headers**

1. Introduction
2. Materials and Methods
  - 2.1. Chemicals
  - 2.2. Animals and tissue preparation
  - 2.3. Competitive substrate/transporter inhibition assay and long-term inhibition
  - 2.4. Data analysis
3. Results
  - 3.1. Method evaluation
  - 3.2. Inhibitory effects by test compounds, dose-effect relationships
  - 3.3. Combinatory effects
  - 3.4. Long-term inhibitory effects
4. Discussion
  - 4.1. MXR efflux inhibition by musks
  - 4.2. Long-term inhibitory effects of musks
  - 4.3. Potential environmental relevance of synthetic musks as chemosensitizers
  - 4.4. Musks and other environmental chemosensitizers – a human health risk?
5. References

## Abstract

Synthetic musk compounds, widely used as fragrances in consumer products, have been detected in human tissue and, surprisingly, in aquatic organisms such as fish and mollusks. Although their persistence and potential to bio-accumulate are of concern, the toxicity and environmental risks of these chemicals are generally regarded as low. Here, however, we show that nitromusks and polycyclic musks inhibit the activity of multi-drug efflux transporters responsible for multi-xenobiotic resistance (MXR) in gills of the marine mussel *Mytilus californianus*. The IC<sub>10</sub> values for the different classes of musks were in the range from 0.09 to 0.39  $\mu$ M and IC<sub>50</sub> values were between 0.74 and 2.56  $\mu$ M. The immediate consequence of inhibition of efflux transporters is that normally excluded xenobiotics will now be able to enter the cell. Remarkably, the inhibitory effects of a brief two-hour exposure to musks were only partially reversed after a 24 to 48-hour recovery period in clean seawater. This unexpected consequence of synthetic musks — a long-term loss of efflux transport activity — will result in continued accumulation of normally excluded toxicants even after direct exposure to the musk has ended. These findings additionally point to the need to determine whether other environmental chemicals have similar long-term effects on these transporters. The results are relevant to human health, as they raise the possibility that exposure to common xenobiotics and pharmaceuticals could cause similar long-term inhibition of these transporters and lead to increased exposure to normally excluded toxicants.